


COMPLETE LISTING OF ALL CLAIMS, WITH MARKINGS AND STATUS IDENTIFIERS
(Currently amended claims showing deletions by ~~strikethrough~~ or [[double brackets]]
and additions by underlining)

1 (currently amended):

A compound of the formula (I),



(I)


the racemic-diastereomeric mixtures and optical isomers of said compound of formula (I) and pharmaceutically-acceptable salts thereof,

wherein

----- represents an optional bond;


R¹ is H, -(CH₂)_m-C(O)-(CH₂)_m-Z¹, -(CH₂)_m-Z¹, -(CH₂)_m-O-Z¹ or -[(C₁-C₆)alkyl]_p-C(O)-NH-(CH₂)_m-Z³;

Z¹ is an optionally substituted moiety selected from the group consisting of (C₁-C₁₂)alkyl, benzo[b]thiophene, phenyl, naphthyl, benzo[b]furanyl, thiophene, isoxazolyl, indolyl,



R² is H or (C₁-C₆)alkyl;

~~or R¹ and R² are taken together with the nitrogen atoms to which they are attached to form a compound of formula (Ia), (Ib) or (Ic),~~



Inventor : Thurieau *et al.*
Serial No. : 10/771,725
Filed : February 4, 2004
Page : 3

R³ is -(CH₂)_m-E-(CH₂)_n-Z²;

E is O, S, C(O), C(O)O, NH C(O)O or a bond;

Z² is H, (C₁-C₁₂)alkyl, amino, (C₁-C₁₂)alkylamino, N,N-di(C₁-C₁₂)alkylamino, (C₁-C₁₂)alkylguanidino, or an optionally substituted moiety selected from the group consisting of phenyl, indolyl, imidazolyl, thiophene, benzothiophene, pyridinyl and naphthyl -CH₂-indol-3-yl, -(CH₂)₄-NH-CO-O-t-Bu or -(CH₂)₄-NH₂;

R⁴ is H or -(CH₂)_m-A¹;

A¹ is -C(=Y)-N(X¹X²), -C(=Y)-X², -C(=NH)-X² or X²;

Y is O or S;


X¹ is H, (C₁-C₁₂)alkyl, -(CH₂)_m-NH-(C₁-C₆)alkyl, -(CH₂)_m-N-di-(C₁-C₆)alkyl or -(CH₂)_m-aryl;


X² is -(CH₂)_m-Y¹-X³ or optionally substituted (C₁-C₁₂)alkyl;

Y¹ is O, S, NH, C=O, (C₂-C₁₂)alkenyl having one or more double bonds, -NH-CO-, -CO-NH-, -NH-CO-O-(CH₂)_m-, -C≡C-, SO₂ or a bond;


X³ is H, an optionally substituted moiety selected from the group consisting of (C₁-C₁₂)alkyl, (C₃-C₈)cycloalkyl, (C₁-C₁₂)alkoxy, aryloxy, (C₁-C₁₂)alkylamino, N,N-di-(C₁-C₁₂)alkylamino, -CH-di-(C₁-C₁₂)alkoxy, pyrrolidinyl, pyridinyl, thiophene, imidazolyl, piperidinyl, piperazinyl, benzothiazolyl, furanyl, indolyl, morpholino, benzo[b]furanyl, quinolinyl, isoquinolinyl, -(CH₂)_m-phenyl, naphthyl, fluorenyl, phthalamidyl, pyrimidinyl,

Chemical structures:





or X^1 and X^2 are taken together with the nitrogen to which they are attached to form an optionally substituted moiety selected from the group consisting of thiazolyl




Y^2 is $CH-X^4$, $N-X^4$, $-C(X^4X^4)$, O or S ;

X^4 for each occurrence is independently $-(CH_2)_m-Y^3-X^5$;

Y^3 is $-C(O)-$, $-C(O)O-$ or a bond;

X^5 is hydroxy, (C_1-C_{12}) alkyl, amino, (C_1-C_{12}) alkylamino, N,N -di- (C_1-C_{12}) alkylamino, or an optionally substituted moiety selected from the group consisting of aryl, aryl(C_1-C_4)alkyl, furanyl, pyridinyl, indolyl, $-CH(phenyl)_2$,




R^5 is $(C_1-C_{12})\text{alkyl}$, $[(C_1-C_6)\text{alkyl}]_m\text{-C(O)-O-}Z^5$, $[(C_1-C_6)\text{alkyl}]_p\text{-C(O)-NH-(CH}_2)_p\text{-}Z^3$ or optionally substituted aryl;

Z^3 for each occurrence is independently amino, $(C_1-C_{12})\text{alkylamino}$, $N,N\text{-di-}(C_1-C_{12})\text{alkylamino}$, $-\text{NH-C(O)-O-(CH}_2)_m\text{-phenyl -NH-C(O)-O-(CH}_2)_m\text{-}(C_1-C_6)\text{alkyl}$ or an optionally substituted moiety selected from the group consisting of imidazolyl, pyridinyl, morpholino, piperidinyl, piperazinyl, pyrazolidinyl, furanyl and thiophene;

R^6 is H or $(C_1-C_6)\text{alkyl}$;

R^7 is $(C_1-C_{12})\text{alkyl}$ or $(CH_2)_m\text{-}Z^4$;

Z^4 is an optionally substituted moiety selected from the group consisting of phenyl, naphthyl, indolyl, thiophene, benzo[b]furan, benzo[b]thiophene, isoxazolyl,



Z^5 is H, $(C_1-C_{12})\text{alkyl}$, $(CH_2)_m\text{-aryl}$;

wherein an optionally substituted moiety is optionally substituted by one or more substituents, each independently selected from the group consisting of Cl, F, Br, I, CF_3 , CN, N_3 , NO_2 , OH, SO_2NH_2 , $-\text{OCF}_3$, $(C_1-C_{12})\text{alkoxy}$, $-(CH_2)_m\text{-phenyl-}(X^6)_n$, $-\text{S-phenyl-}(X^6)_n$, $-\text{S-}(C_1-C_{12})\text{alkyl}$, $-\text{O-(CH}_2)_m\text{-phenyl-}(X^6)_n$, $-(CH_2)_m\text{-C(O)-O-(C}_1\text{-C}_6)\text{alkyl}$, $-(CH_2)_m\text{-C(O)-(C}_1\text{-C}_6)\text{alkyl}$, $-\text{O-(CH}_2)_m\text{-NH}_2$, $-\text{O-(CH}_2)_m\text{-NH-}(C_1-C_6)\text{alkyl}$, $-\text{O-(CH}_2)_m\text{-N-di-((C}_1\text{-C}_6)\text{alkyl)}$ and $-[(C_1-C_{12})\text{alkyl}]_p\text{-}(X^6)_n$;

X^6 for each occurrence is independently selected from the group consisting of hydrogen, Cl, F, Br, I, NO_2 , N_3 , CN, OH, $-\text{CF}_3$, $-\text{OCF}_3$, $(C_1-C_{12})\text{alkyl}$, $(C_1-C_{12})\text{alkoxy}$, $-(CH_2)_m\text{-NH}_2$, $-(CH_2)_m\text{-NH -(C}_1\text{-C}_6)\text{alkyl}$, $-(CH_2)_m\text{-N-di-((C}_1\text{-C}_6)\text{alkyl)}$ and $-(CH_2)_m\text{-phenyl}$;

m for each occurrence is independently 0 or an integer from 1 to 6;

n for each occurrence is independently an integer from 1 to 5; and

p for each occurrence is independently 0 or 1[[:]]

Inventor : Thurieau *et al.*
Serial No. : 10/771,725
Filed : February 4, 2004
Page : 6

~~provided that:~~

- (a) when R^5 is (C_1-C_{12}) alkyl, or $C(O)OZ^5$ and Z^5 is (C_1-C_{12}) alkyl or optionally substituted aryl; R^6 is H or (C_1-C_6) alkyl; R^7 is (C_1-C_{12}) alkyl or Z^4 and Z^4 is thiophene or optionally substituted phenyl, then R^3 is not $C(O)O(CH_2)_mZ$ where m is 0 and Z is H or (C_1-C_{12}) alkyl or where m is 1 to 6 and Z is H;
- (b) when R^5 is (C_1-C_{12}) alkyl or optionally substituted phenyl; R^6 is H or (C_1-C_6) alkyl; R^7 is (C_1-C_{12}) alkyl and R^3 is $O(CH_2)Z^2$, then Z^2 is not an optionally substituted moiety selected from the group consisting of phenyl, indolyl, imidazolyl, thiophene, benzothiophene, pyridinyl, and naphthyl;
- (c) when R^5 is (C_1-C_{12}) alkyl; R^6 is (C_1-C_6) alkyl; R^7 is (C_1-C_{12}) alkyl; and R^3 is OZ^2 or SZ^2 , then Z^2 is not an optionally substituted moiety selected from the group consisting of phenyl, naphthyl, thiophene, benzothienyl and indolyl[.]; and
- (d) at least one m in the definition of R^3 must be an integer from 1 to 6.

2 (canceled)

3 (canceled)

4 (canceled)


5 (withdrawn): A compound according to claim 1 wherein R^1 is H; R^2 is H; R^3 is $-CH_2\text{-indol-3-yl}$; R^4 is $-(CH_2)_m-A^1$ where m in the definition of R^4 is 0; R^5 is phenyl or t-Bu; R^6 is H;

A^1 is $-C(=Y)-N(X^1X^2)$;

Y is O or S; X^1 is H; X^2 is $-(CH_2)_m-Y^1-X^3$;

m in the definition of X^2 is 0, 1 or 2;

Y^1 is a bond; and X^3 is phenyl, o-Cl-phenyl, m-Cl-phenyl, p-phenyloxy-phenyl, 2,6-di-isopropylphenyl, m-CF₃-phenyl, p-ethoxycarbonyl-phenyl, 2,4-difluorophenyl, m-NO₂-phenyl, p-benzyloxyphenyl, o-isopropylphenyl, n-hexyl, 4-morpholino, naphthyl or



Inventor : Thurieau *et al.*
Serial No. : 10/771,725
Filed : February 4, 2004
Page : 7


6 (withdrawn): A compound according to claim 1 wherein R¹ is H; R² is H; R³ is -CH₂-indol-3-yl; R⁴ is -(CH₂)_m-A¹ where m in the definition of R⁴ is 0; R⁵ is phenyl or t-Bu; R⁶ is H;

where A¹ is -C(=Y)-X²;

Y is O; X² is -(CH₂)_m-Y¹-X³;

where m in the definition of X² is 0, 1 or 2;

Y¹ is O, -CO-NH-, -NH-CO-O-CH₂-or a bond; and X³ is methyl, 3-pentyl, phenyl, p-NO₂-phenyl, phthalamidyl, N,N-dimethylamino, p-aminophenyl, fluorenyl or




7 (withdrawn): A compound according to claim 1 wherein R¹ is H; R² is H; R³ is -CH₂-indol-3-yl; R⁴ is -(CH₂)_m-A¹ where m in the definition of R⁴ is 0; R⁵ is phenyl or t-Bu; R⁶ is H;

where A¹ is -C(=Y)-N(X¹X²);

Y is O; X¹ is hydrogen; X² is -(CH₂)_m-Y¹-X³;

where m in the definition of X² is 0, 1, 2 or 3;

Y¹ is O, or a bond; and X³ is cyclopentyl, 4-OH-butyl, N,N-diethylamino, N-methyl-pyrrolidin-3-yl, -CH(ethoxy)₂, phenyl, p-SO₂NH₂-phenyl p-OH-phenyl, o-CF₃-phenyl, p-Cl-phenyl, -CH(phenyl)₂,



8 (withdrawn): A compound according to claim 1 wherein R¹ is H; R² is H; R³ is -CH₂-indol-3-yl; R⁴ is -(CH₂)_m-A¹ where m in the definition of R⁴ is 0; R⁵ is phenyl or t-Bu; R⁶ is H;

Inventor : Thurieau *et al.*
Serial No. : 10/771,725
Filed : February 4, 2004
Page : 8

where A^1 is $-C(=Y)-X^2$;

Y is O; X^2 is $-(CH_2)_m-Y^1-X^3$;

where m in the definition of X^2 is 0, 1, 2 or 3;

Y^1 is $-NH-CO$, $-C=C-$, $-C\equiv C-$ or a bond; and X^3 is t-butyl, 1-methylcarbonyl-piperidin-4-yl, phenyl, p-Cl-phenyl, m-CF₃-phenyl, 4-nitro-naphthyl, p-methoxy-phenyl, m-(phenylethyl)-phenyl, indol-3-yl or p-aminophenyl.

9 (currently amended): A compound according to claim 1 wherein R^1 is H;

R^2 is H; R^3 is ~~CH₂-indol-3-yl, (CH₂)₄-NH-CO-O-t-Bu or (CH₂)₄-NH₂~~; R^5 is phenyl, o-methoxyphenyl, p-Br-phenyl, p-nitro-phenyl or p-N,N-diethylamino-phenyl; R^6 is H.

10 (withdrawn): A compound according to claim 1 wherein R^1 is H; R^2 is H;

R^3 is ~~CH₂-indol-3-yl, (CH₂)₄-NH-CO-O-t-Bu or (CH₂)₄-NH₂~~; R^4 is $-(CH_2)_m-A^1$ where m in the definition of R^4 is 0; R^5 is phenyl, o-methoxyphenyl, p-methoxyphenyl, p-Br-phenyl, p-nitro-phenyl or p-N,N-diethylamino-phenyl; R^6 is H;

where A^1 is $-C(=Y)-X^2$;

Y is O; X^2 is $-(CH_2)_m-Y^1-X^3$;

where m in the definition of X^2 is 1;

Y^1 is a bond; and X^3 is phenyl, o-Br-phenyl, m-Br-phenyl, p-Br-phenyl, o-Cl-phenyl, m-Cl-phenyl, p-Cl-phenyl, o-nitro-phenyl, m-nitro-phenyl, p-nitro-phenyl, o-CF₃-phenyl, m-CF₃-phenyl, p-CF₃-phenyl, o-F-phenyl, m-F-phenyl, p-F-phenyl, N,N-di-methylamino-phenyl, o-OMe-phenyl, m-OMe-phenyl, p-OMe-phenyl, 3,4-di-Cl-phenyl, 3,4,5-tri-OMe-phenyl, p-Me-phenyl, p-OH-phenyl or 2,4-di-F-phenyl.

11 (original): A compound according to claim 9 wherein R^5 is phenyl and R^3 is

$-(CH_2)$ -indol-3-yl and the stereochemistry at the carbon to which R^3 is attached is the R-configuration.

Inventor : Thurieau *et al.*
Serial No. : 10/771,725
Filed : February 4, 2004
Page : 9

12 (withdrawn): A compound according to claim 10 wherein R⁵ is phenyl and R³ is -(CH₂)-indol-3-yl and the stereochemistry at the carbon to which R³ is attached is the R-configuration.

13 (withdrawn): A compound according to claim 10 wherein R⁵ is o-OMe-phenyl and R³ is -(CH₂)-indol-3-yl and the stereochemistry at the carbon to which R³ is attached is the R-configuration.

14 (withdrawn): A compound according to claim 10 wherein R⁵ is o-OMe-phenyl and R³ is -(CH₂)-indol-3-yl and the stereochemistry at the carbon to which R³ is attached is the S-configuration.

15 (withdrawn): A compound according to claim 1 wherein R¹ is H; R² is H; R³ is -(CH₂)₄-NH-CO-O-t-Bu or -(CH₂)₄-NH₂; R⁴ is -(CH₂)_m-A¹ where m in the definition of R⁴ is 0; R⁵ is phenyl; R⁶ is H;

where A¹ is -C(=Y)-X²;

Y is O; X² is -(CH₂)_m-Y¹-X³;

where m in the definition of X² is 0, 1 or 2;

Y¹ is S, SO₂ or a bond; and X³ is phenyl, 3,4-di-Cl-phenyl, 3,4,5-tri-OMe-phenyl, p-Me-phenyl, p-OH-phenyl, 2,4-di-F-phenyl, 2-furanyl, 2-pyridinyl, 3-pyridinyl, naphthyl, 2-quinolinyl, 3-quinolinyl, 4-quinolinyl, 8-quinolinyl, 1-isoquinolinyl, 2-thiophene or 2-pyrimidinyl.

16 (withdrawn): A compound according to claim 1 wherein R¹ is H; R² is H; R³ is -(CH₂)₄-NH-CO-O-t-Bu or -(CH₂)₄-NH₂; R⁴ is -(CH₂)_m-A¹ where m in the definition of R⁴ is 0; R⁵ is phenyl; R⁶ is H;

where A¹ is -C(=Y)-X²;

Y is O; X² is -(CH₂)_m-Y¹-X³;

where m in the definition of X² is 0, 1, 2 or 3;

Inventor : Thurieau *et al.*
Serial No. : 10/771,725
Filed : February 4, 2004
Page : 10

Y^1 is a bond; and X^3 is 5-indolyl, 3-indolyl, 4-indolyl, 2-indolyl, 5-OMe-indol-3-yl, 5-OMe-indol-2-yl, 5-OH-indol-2-yl, 5-OH-indol-3-yl, 5-Br-indol-3-yl, 2-Me-indol-3-yl, 2-benzothiophene, 3-benzothiophene or 2-benzofuran.

17 (withdrawn): A compound according to claim 1 wherein R^1 is H; R^2 is H; R^3 is $-(CH_2)_m\text{-indol-3-yl}$, $-(CH_2)_4\text{NH-CO-O-t-Bu}$ or $-(CH_2)_4\text{NH}_2$; R^4 is $-(CH_2)_m\text{-A}^1$ where m in the definition of R^4 is 0; R^5 is phenyl, o-OMe-phenyl or p-OMe-phenyl; R^6 is H; where A^1 is X^2 ;

X^2 is $-(CH_2)_m\text{-Y}^1\text{-X}^3$;

where m in the definition of X^2 is 1, 2 or 3;

Y^1 is S, O or a bond; and X^3 is phenyl, o-OH-phenyl, p-OH-phenyl, o-F-phenyl, m-F-phenyl, p-F-phenyl, o-CF₃-phenyl, o-OMe-phenyl, m-OMe-phenyl, o-nitro-phenyl, p-nitro-phenyl, 3,4-di-Cl-phenyl, 2-nitro-3-OMe-phenyl, o-Br-phenyl, m-Br-phenyl, p-Br-phenyl, 2-thiophene, 3,4,5-tri-OMe-phenyl, p-N,N-dimethylamino-phenyl, p-OCF₃-phenyl, p-(3-(N,N-dimethylamino)propoxy)phenyl, 3-F-4-OMe-phenyl, 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, 2-Cl-quinolin-3-yl, 2-quinolinly, methyl, n-butyl, n-pentyl, n-hexyl, 3,3-dimethyl-butyl, benzyl, cyclohexyl or p-t-Bu-phenyl.

18 (withdrawn): A compound according to claim 1 wherein R^1 is H; R^2 is H; R^3 is $-(CH_2)_4\text{-NH-CO-O-t-Bu}$ or $-(CH_2)_4\text{-NH}_2$; R^4 is $-(CH_2)_m\text{-A}^1$ where m in the definition of R^4 is 0; R^5 is phenyl; R^6 is H;

where A^1 is X^2 ;

X^2 is $-(CH_2)_m\text{-Y}^1\text{-X}^3$;

where m in the definition of X^2 is 1, 2 or 3;


Y^1 is O or a bond; and X^3 is phenyl, o-OH-phenyl, p-OH-phenyl, o-F-phenyl, m-F-phenyl, p-F-phenyl, o-CF₃-phenyl, o-OMe-phenyl, m-OMe-phenyl, p-OMe-phenyl, o-nitro-phenyl, p-nitro-phenyl, 3,4-di-Cl-phenyl, 2-nitro-3-OMe-phenyl, o-Br-phenyl, m-Br-phenyl, p-Br-phenyl, p-phenyl-phenyl, 2-thiophene, 3,4,5-tri-OMe-phenyl, p-N,N-dimethylamino-phenyl, p-benzyloxy-phenyl, p-OCF₃-phenyl, p-(3-(N,N-dimethylamino)propoxy)phenyl, 3-F-4-OMe-phenyl, 2-pyridinyl, 3-pyridinyl, 4-pyridinyl,

Inventor : Thurieau *et al.*
Serial No. : 10/771,725
Filed : February 4, 2004
Page : 11

2-Cl-quinolin-3-yl, 2-quinolinly, 3-indolyl, 6-methoxycarbonyl-indol-3-yl, 1-methyl-indol-3-yl, 2-methyl-indol-3-yl, methyl, n-butyl, n-pentyl, n-hexyl, 3,3-dimethyl-butyl, benzyl, cyclohexyl or p-t-Bu-phenyl.

19 (withdrawn): A compound according to claim 1 wherein R¹ is -(CH₂)-CO-Z¹; R² is H; R³ is -(CH₂)₄-NH-CO-O-t-Bu, ~~(CH₂)₄NH-CO-O-benzyl~~, ~~(CH₂)-phenyl~~ or -(CH₂)-indol-3-yl; R⁴ is -(CH₂)_m-A¹ where m in the definition of R⁴ is 0; R⁵ is phenyl; R⁶ is H; where Z¹ is ethyl, phenyl, p-OMe-phenyl, p-phenyl-phenyl, p-Cl-phenyl, p-Br-phenyl, p-N₃-phenyl, p-F-phenyl, m-nitro-phenyl, p-nitro-phenyl, p-CN-phenyl, 2,5-di-OMe-phenyl, 3,4-di-Cl-phenyl, N,N-dimethylamino-phenyl, 3-methyl-4-Cl-phenyl or naphthyl; A¹ is -C(=Y)-X²;
Y is O; X² is -(CH₂)_m-Y¹-X³;
where m in the definition of X² is 0;
Y¹ is O; and X³ is t-Bu.

20 (withdrawn): A compound according to claim 1 wherein R¹ is -(CH₂)-CO-(CH₂)_m-Z¹ where m in the definition of R¹ is 0, 1 or 2; R² is H; R³ is -(CH₂)-indol-3-yl or -(CH₂)₄-NH-CO-O-t-Bu; R⁴ is H or -(CH₂)_m-A¹ where m in the definition of R⁴ is 0; R⁵ is phenyl, o-OMe-phenyl, p-nitro-phenyl, p-Br-phenyl, t-Bu, -CH(CH₃)₂-CO-NH-(CH₂)₂-CO-O-t-Bu, -CH(CH₃)₂-CO-NH-(CH₂)₃-imidazol-1-yl, -CH(CH₃)₂-CO-NH-(CH₂)₂-pyridin-2-yl, -CH(CH₃)₂-CO-NH-(CH₂)₃-4-morpholino, -CH(CH₃)₂-CO-NH-(CH₂)-pyridin-4-yl or -CH(CH₃)₂-CO-NH-(CH₂)₂-N,N-diethylamino; R⁶ is H;
where Z¹ is ethyl, propyl, phenyl, p-OMe-phenyl, p-Cl-phenyl, p-Br-phenyl, p-F-phenyl, p-nitro-phenyl, m-nitro-phenyl, p-CN-phenyl, p-N₃-phenyl, p-phenyl-phenyl, 3-Me-4-Cl-phenyl, p-N,N-diethylamino-phenyl, 2,5-di-OMe-phenyl, 3,4-di-Cl-phenyl, 3,4-di-F-phenyl, p-OCF₃-phenyl, p-benzyloxy-phenyl, p-pentyl-phenyl, 3,4,5-tri-OMe-phenyl, 3-nitro-4-Cl-phenyl, 3-Cl-4-nitro-phenyl, 3-methyl-5-chloro-benzothiophen-2-yl, 2-benzofuranyl, 3-benzothiophene, 3-phenyl-isoxazol-5-yl, 3-(2,4-di-Cl-phenyl)-isoxazol-5-yl, 3-indolyl, 5-Br-thiophen-2-yl, naphthyl,



A¹ is -C(=Y)-X²;

Y is O; X² is -(CH₂)_m-Y¹-X³;

where m in the definition of X² is 0;

Y¹ is O; and X³ is t-Bu.

21 (canceled)

22 (canceled)

23 (canceled)

24 (canceled)

25 (canceled)

26 (canceled)

27 (canceled)

28 (canceled)

29 (canceled)

30 (withdrawn): A pharmaceutical composition comprising a compound according to claim 1 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

31 (withdrawn): A method of eliciting an agonist effect from one or more of a somatostatin subtype receptor in a subject in need thereof, which comprises administering a compound according to claim 1 or a pharmaceutically acceptable salt thereof to said subject.

32 (withdrawn): A method of eliciting an antagonist effect from one or more of a somatostatin subtype receptor in a subject in need thereof, which comprises administering a compound according to claim 1 or a pharmaceutically acceptable salt thereof to said subject.

33 (withdrawn): A method of binding one or more of a somatostatin subtype receptor in a subject in need thereof, which comprises administering a compound according to claim 1 or a pharmaceutically acceptable salt thereof to said subject.

Inventor : Thurieau *et al.*
Serial No. : 10/771,725
Filed : February 4, 2004
Page : 13

34 (withdrawn): A method of treating acromegaly, restenosis, Crohn's disease, systemic sclerosis, external and internal pancreatic pseudocysts and ascites, VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, Zollinger-Ellison Syndrome, diarrhea, AIDS related diarrhea, chemotherapy related diarrhea, scleroderma, Irritable Bowel Syndrome, pancreatitis, small bowel obstruction, gastroesophageal reflux, duodenogastric reflux, Cushing's Syndrome, gonadotropinoma, hyperparathyroidism, Graves' Disease, diabetic neuropathy, Paget's disease, polycystic ovary disease, cancer, cancer cachexia, hypotension, postprandial hypotension, panic attacks, GH secreting adenomas or TSH secreting adenomas, in a subject in need thereof, which comprises administering a compound according to claim 1 or a pharmaceutically acceptable salt thereof to said subject.

35 (withdrawn): A method of treating diabetes mellitus, hyperlipidemia, insulin insensitivity, Syndrome X, angiopathy, proliferative retinopathy, dawn phenomenon, Nephropathy, peptic ulcers, enterocutaneous and pancreaticocutaneous fistula, Dumping syndrome, watery diarrhea syndrome, acute or chronic pancreatitis, gastrointestinal hormone secreting tumors, angiogenesis, inflammatory disorders, chronic allograft rejection, angioplasty, graft vessel bleeding or gastrointestinal bleeding in a subject in need thereof, which comprises administering a compound according to claim 1 or a pharmaceutically acceptable salt thereof to said subject.

36 (withdrawn): A method of inhibiting the proliferation of helicobacter pylori in a subject in need thereof, which comprises administering a compound according claim 1 or a pharmaceutically acceptable salt thereof, to said subject.

37 (canceled)

38 (canceled)

39 (canceled)

40 (canceled)

41 (canceled)

42 (canceled)

43 (canceled)